Last Updated: April 29, 2002

Reported[1] cases and rates of notifiable STD from January 1 to December 31, 2001
and January 1 to December 31, 2000

**Division of Sexual Health Promotion and STD Prevention & Control, Centre for Infectious Disease Prevention and Control, Health Canada**

| Province/ Territory | Genital Chlamydia | | | | Gonorrhea | | | | Infectious syphilis[4] | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | National Goal: 50 per 100,000 by 2010 | | | | National Goal: Elimination by 2010 | | | | National Goal: <0.5 per 100,000 | | | |
| | Year to Date Cases | | 2000 Rate[2] | 2001 Rate projected[3] | Year to Date Cases | | 2000 Rate[2] | 2001 Rate projected[3] | Year to Date Cases | | 2000 Rate[2] | 2001 Rate projected[3] |
| | 2000 | 2001 | | | 2000 | 2001 | | | 2000 | 2001 | | |
| Canada* | 46397 | 48683 | 150.9 | 159.4 | 6225 | 6478 | 20.2 | 21.2 | 174 | 280 | 0.6 | 0.9 |
| BC | 6193 | 5792 | 152.7 | 140.7 | 724 | 612 | 17.9 | 14.9 | 95 | 176 | 2.3 | 4.3 |
| AB | 6001 | 6462 | 202.4 | 213.3 | 586 | 801 | 19.5 | 26.4 | 13 | 22 | 0.4 | 0.7 |
| SK* | 2921 | 2204 | 284.2 | 286.8 | 461 | 329 | 44.7 | 42.8 | 1 | 2 | 0.1 | 0.2 |
| MB | 3263 | 3254 | 284.0 | 283.0 | 658 | 700 | 57.3 | 60.9 | 1 | 0 | 0.1 | 0.0 |
| ON | 14660 | 16114 | 125.8 | 136.4 | 2819 | 2885 | 24.2 | 24.4 | 45 | 45 | 0.4 | 0.4 |
| QC | 8676 | 10088 | 117.8 | 136.5 | 670 | 826 | 9.1 | 11.2 | 7 | 14 | 0.1 | 0.2 |
| NB | 1157 | 1138 | 153.1 | 150.3 | 10 | 9 | 1.3 | 1.2 | 1 | 0 | 0.1 | 0.0 |
| NS | 1406 | 1621 | 149.1 | 171.9 | 57 | 86 | 6.0 | 9.1 | 1 | 0 | 0.1 | 0.0 |
| NF | 554 | 597 | 102.9 | 111.2 | 5 | 0 | 0.9 | 0.0 | 0 | 1 | 0.0 | 0.2 |
| PE | 231 | 150 | 166.3 | 107.6 | 0 | 0 | 0.0 | 0.0 | 0 | 0 | 0.0 | 0.0 |
| YK | 144 | 129 | 485.8 | 421.5 | 5 | 3 | 16.9 | 9.8 | 10 | 20 | 33.7 | 65.3 |
| NW | 487 | 533 | 1150.1 | 1259.6 | 135 | 151 | 318.8 | 356.8 | 0 | 0 | 0.0 | 0.0 |
| NN | 704 | 601 | 2551.5 | 2129.5 | 95 | 76 | 346.8 | 269.3 | 0 | 0 | 0.0 | 0.0 |

[1] Data are preliminary and expected to change: 2001 data expected to change more than 2000 data.

[2] Rate per 100,000 population based on reported cases for 2000. Population estimates provided by Statistics Canada.

[3] Rate is calculated by projecting current number of cases to 12 months, then calculating the rate (per 100,000) using 2001 population estimates provided by Statistics Canada. No adjustment has been made for seasonal variability or reporting delays.

[4] Infectious syphilis: primary, secondary and early latent syphilis.

*2001 data of Canada, SK are incomplete and the rates are projected to the end of 2001 using the available data

**Source**: Division of Sexual Health Promotion and STD Prevention & Control, Centre for Infectious Disease Prevention and Control, Health Canada

Last Updated: April 29, 2002

Reported[1] cases and rates of notifiable STD from January 1 to December 31, 2001
and January 1 to December 31, 2000

**Division of Sexual Health Promotion and STD Prevention & Control, Centre for Infectious Disease Prevention and Control, Health Canada**

| Province/ Territory | Genital Chlamydia | | | | Gonorrhea | | | | Infectious syphilis[4] | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | National Goal: 50 per 100,000 by 2010 | | | | National Goal: Elimination by 2010 | | | | National Goal: <0.5 per 100,000 | | | |
| | Year to Date Cases | | 2000 Rate[2] | 2001 Rate projected[3] | Year to Date Cases | | 2000 Rate[2] | 2001 Rate projected[3] | Year to Date Cases | | 2000 Rate[2] | 2001 Rate projected[3] |
| | 2000 | 2001 | | | 2000 | 2001 | | | 2000 | 2001 | | |
| **Canada*** | **46397** | **48683** | **150.9** | **159.4** | **6225** | **6478** | **20.2** | **21.2** | **174** | **280** | **0.6** | **0.9** |
| BC | 6193 | 5792 | 152.7 | 140.7 | 724 | 612 | 17.9 | 14.9 | 95 | 176 | 2.3 | 4.3 |
| AB | 6001 | 6462 | 202.4 | 213.3 | 586 | 801 | 19.5 | 26.4 | 13 | 22 | 0.4 | 0.7 |
| SK* | 2921 | 2204 | 284.2 | 286.8 | 461 | 329 | 44.7 | 42.8 | 1 | 2 | 0.1 | 0.2 |
| MB | 3263 | 3254 | 284.0 | 283.0 | 658 | 700 | 57.3 | 60.9 | 1 | 0 | 0.1 | 0.0 |
| ON | 14660 | 16114 | 125.8 | 136.4 | 2819 | 2885 | 24.2 | 24.4 | 45 | 45 | 0.4 | 0.4 |
| QC | 8676 | 10088 | 117.8 | 136.5 | 670 | 826 | 9.1 | 11.2 | 7 | 14 | 0.1 | 0.2 |
| NB | 1157 | 1138 | 153.1 | 150.3 | 10 | 9 | 1.3 | 1.2 | 1 | 0 | 0.1 | 0.0 |
| NS | 1406 | 1621 | 149.1 | 171.9 | 57 | 86 | 6.0 | 9.1 | 1 | 0 | 0.1 | 0.0 |
| NF | 554 | 597 | 102.9 | 111.2 | 5 | 0 | 0.9 | 0.0 | 0 | 1 | 0.0 | 0.2 |
| PE | 231 | 150 | 166.3 | 107.6 | 0 | 0 | 0.0 | 0.0 | 0 | 0 | 0.0 | 0.0 |
| YK | 144 | 129 | 485.8 | 421.5 | 5 | 3 | 16.9 | 9.8 | 10 | 20 | 33.7 | 65.3 |
| NW | 487 | 533 | 1150.1 | 1259.6 | 135 | 151 | 318.8 | 356.8 | 0 | 0 | 0.0 | 0.0 |
| NN | 704 | 601 | 2551.5 | 2129.5 | 95 | 76 | 346.8 | 269.3 | 0 | 0 | 0.0 | 0.0 |

[1] Data are preliminary and expected to change: 2001 data expected to change more than 2000 data.

[2] Rate per 100,000 population based on reported cases for 2000. Population estimates provided by Statistics Canada.

[3] Rate is calculated by projecting current number of cases to 12 months, then calculating the rate (per 100,000) using 2001 population estimates provided by Statistics Canada. No adjustment has been made for seasonal variability or reporting delays.

[4] Infectious syphilis: primary, secondary and early latent syphilis.

*2001 data of Canada, SK are incomplete and the rates are projected to the end of 2001 using the available data

**Source**: Division of Sexual Health Promotion and STD Prevention & Control, Centre for Infectious Disease Prevention and Control, Health Canada

**Cumulative cases of genital chlamydia from January 1 to December 31, 2001 and January 1 to December 31, 2000**



**Cumulative cases of gonorrhea from January 1 to December 31, 2001 and January 1 to December 31, 2000**



## Reported rate of genital chlamydia in 2000 and projected rate for 2001



## Reported rate of Gonorrhea in 2000 and projected rate for 2001



IMPORTANT NOTE CONCERNING THE FOLLOWING PAGES

THE PAGES WHICH FOLLOW HAVE BEEN FILMED TWICE IN ORDER TO OBTAIN THE BEST REPRODUCTIVE QUALITY

USERS SHOULD CONSULT ALL THE PAGES REPRODUCED ON THE FICHE IN ORDER TO OBTAIN A COMPLETE READING OF THE TEXT.

REMARQUE IMPORTANTE CONCERNANT LES PAGES QUI SUIVENT

LES PAGES SUIVANTES ONT ÉTÉ REPRODUITES EN DOUBLE AFIN D'AMÉLIORER LA QUALITÉ DE REPRODUCTION

LES UTILISATEURS DOIVENT CONSULTER TOUTES LES PAGES REPRODUITES SUR LA FICHE AFIN D'OBTENIR LA LECTURE DU TEXTE INTÉGRAL

Micromedia
ProQuest

[illegible]

# Comparaison des cas rapportés[1] et des taux de MTS à déclaration obligatoire du 1er janvier au 31 décembre 2001 et du 1er janvier au 31 décembre 2000

## Division de la promotion de la santé sexuelle, de la prévention et du contrôle des MTS, Centre de prévention et de contrôle des maladies infectieuses, Santé Canada

| Province/territoire | Chlamydiose génitale — déclaré facultatif — nb de cas — taux/100 000 pers. (2001 2000) | | | | Gonorrhée — déclaré national — nombre de cas, 2001 2000 | | | | Syphilis infectieuse[2] — déclaré national — nbre de cas et taux/100 000 pers. | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | nbre de cas 2001 | nbre de cas 2000 | taux 2001 | taux prov. 2001 | nbre de cas 2001 | nbre de cas 2000 | taux 2001 | taux prov. 2001 | nbre de cas 2001 | nbre de cas 2000 | taux 2001 | taux prov. 2001 |
| Canada[3] | 45 107 | 43 [illegible] | 149.0 | 149.0 | 8 224 | 8 [illegible] | 27.0 | 27.0 | 1 [illegible] | [illegible] | 3.0 | 3.0 |
| T.-N. | 1 [illegible] | 1 [illegible] | 192.7 | 192.7 | 1 [illegible] | 1 [illegible] | 17.0 | 19.0 | 80 | 1 [illegible] | 6.0 | 9.0 |
| Î.-P.-É. | 1 [illegible] | 1 [illegible] | 116.0 | 116.0 | [illegible] | [illegible] | 13.0 | 20.0 | 13 | 20 | 9.0 | 14.0 |
| Nouv.[4] | 2 [illegible] | 2 [illegible] | 286.0 | 286.0 | [illegible] | [illegible] | 99.7 | 92.0 | 1 | 2 | 0.1 | 0.3 |
| Yukon | [illegible] | [illegible] | [illegible] | [illegible] | [illegible] | [illegible] | 47.0 | 83.0 | 1 | 1 | 0.1 | 0.1 |
| Ont. | 1 [illegible] | 1 [illegible] | 152.0 | 148.0 | [illegible] | [illegible] | 30.0 | 30.0 | 1 [illegible] | 1 [illegible] | 1.0 | 1.0 |
| Qc | [illegible] | [illegible] | 117.0 | 148.0 | [illegible] | [illegible] | 3.1 | 10.2 | 7 | 14 | 0.1 | 0.2 |
| N.-B. | 1 [illegible] | 1 [illegible] | 150.0 | 150.0 | 15 | 9 | 1.3 | 1.5 | 1 | 0 | 0.1 | 0.0 |
| N.-É. | 1 [illegible] | 1 [illegible] | 143.0 | 175.0 | 27 | 37 | 3.0 | 3.1 | 1 | 1 | 0.1 | 0.1 |
| T.-N. | 2 [illegible] | 2 [illegible] | 142.0 | [illegible] | 2 | 1 | 1.0 | 1.0 | 0 | 1 | 0.0 | 0.2 |
| Î.-P.-É. | [illegible] | 1 [illegible] | 100.0 | 107.0 | 1 | 1 | 1.0 | 1.0 | 1 | 1 | 0.0 | 0.0 |
| Yukon | 1 [illegible] | 1 [illegible] | 385.0 | 401.2 | 2 | 3 | 15.0 | 5.0 | 15 | 22 | 55.7 | 82.0 |
| T.N.-O. | 1 [illegible] | 2 [illegible] | 1 [illegible] | 1 [illegible] | 1 [illegible] | 1 [illegible] | 3 [illegible] | 3 [illegible] | 1 | 1 | 0.0 | 0.0 |
| Nunavut | 1 [illegible] | 1 [illegible] | 2 [illegible] | 2 [illegible] | 2 [illegible] | 1 [illegible] | 3 [illegible] | 2 [illegible] | 0 | 0 | 0.0 | 0.0 |

[illegible] cas déclarés sont préliminaires et les chiffres ci-dessus peuvent changer. Les données de 2001 sont définitives lorsque plus que cette date. [illegible]

[illegible] taux pour 100 000 personnes ci-joint des cas rapportés pour 2000. Les estimations démographiques provisoires de Statistique Canada.

[illegible] Ils ont permis de faire entre les taux fondés sur population de certains grand de ce cas. Par ex., pour calculer calculé à taux pour 100 000 personnes ci-joint des données démographiques de 2000 de Statistique Canada. Ils n'ont pas tenu compte des variations saisonnières ou des écarts dans les rapports.

[illegible] La syphilis infectieuse : la syphilis primaire et secondaire et latente précoce.

[illegible] Les données de 2001 sont [illegible]. Elles sont incomplètes à cet égard sont projetés pour à fin de 2001 en [illegible] les données disponibles.

Division de la promotion de la santé sexuelle, de la prévention et du contrôle des MTS, Centre de prévention et de contrôle des maladies infectieuses

[illegible]

[illegible]

[illegible]

| [illegible] | [illegible] | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| [illegible] | [illegible] | | | | | | | | | | | | |
| [illegible] | [illegible] | | | | | | | | | | | | |
| [illegible] | [illegible] | | | | | | | | | | | | |
| [illegible] | [illegible] | | | | | | | | | | | | |
| [illegible] | [illegible] | | | | | | | | | | | | |
| [illegible] | [illegible] | | | | | | | | | | | | |
| [illegible] | [illegible] | | | | | | | | | | | | |
| [illegible] | [illegible] | | | | | | | | | | | | |
| [illegible] | [illegible] | | | | | | | | | | | | |
| [illegible] | [illegible] | | | | | | | | | | | | |
| [illegible] | [illegible] | | | | | | | | | | | | |
| [illegible] | [illegible] | | | | | | | | | | | | |
| [illegible] | [illegible] | | | | | | | | | | | | |
| [illegible] | [illegible] | | | | | | | | | | | | |
| [illegible] | [illegible] | | | | | | | | | | | | |

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

**Total des cas de chlamydiose génitale du 1er janvier au 31 décembre 2001, et du 1er janvier au 31 décembre, 2000**



**Total des cas de gonorrhée du 1er janvier au 31 décembre 2001, et du 1er janvier au 31 décembre, 2000**



## Taux rapporté de chlamydiose génitale en 2000 et taux prévu pour 2001



## Taux rapporté de gonorrhée en 2000 et taux prévu pour 2001